

— ABLYBIO, Help Your Research



CD5L Rabbit pAb

货号: **AYP16819**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	38kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse liver
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-347 of human CD5L (NP_005885.1).
------	---

靶点信息

研究背景	CD5L (CD5 antigen-like), also known as Sp alpha and AIM, is a 50 kDa secreted glycoprotein that belongs to the SRCR (scavenger receptor cysteine rich) group B family of proteins. CD5L is up-regulated in macrophages at inflammatory sites. It sustains inflammatory reactions by both increasing the phagocytic capacity of macrophages and impeding the apoptosis of local macrophages, NK cells, and T cells. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gram negative bacteria. In the thymic cortex, CD5L protects cortical CD4+CD8+thymocytes from apoptosis. CD5L circulates in the serum in complex with IgM.
基因ID	922
基因名	CD5L
Swiss	O43866 (https://www.uniprot.org/uniprotkb/O43866/entry)
别名	CD5L,AIM,API6,CT-2,PRO229,SP-ALPHA,Spalpha,hAIM,CD5L Rabbit pAb,Apoptosis inhibitor expressed by macrophages,IgM-associated peptide

产品验证



Western blot analysis of CD5L expressed in Raji, Mouse liver using CD5L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)