

TCF19 Rabbit pAb

货号: **AYP16836**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	37kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse thymus,Mouse spleen,Mouse brain,Rat spleen
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-290 of human TCF19 (NP_009040.2).
序列	MLPCFQLLRIGGGRRGGDLYTFHPPAGAGCTYRLGHRADLCDVALRPQQEPLISGIHAELHAEPRGDDWRVSLSDHSSQ GTLVNNVRLPRGHRLELSDGDLLTFGPEGPPGTSPSEFYFMFQQVRVKPQDFAAITIPRSRGEARVGAGFRPMLPSQGAP QRPLSTFSPAPKATLILNSIGLSKLRPQPLTFSPSWGPKSLPVPAPPGEMGTTPSAPPQRNRRKSVHRVLAELDDESEPP ENPPPVLMERKLRVDKAPLPTGNRRGRPRKYPVSAPMAPPVAVGGGE

靶点信息

研究背景	This gene encodes a protein that contains a PHD-type zinc finger domain and likely functions as a transcription factor. The encoded protein plays a role proliferation and apoptosis of pancreatic beta cells. Alternative splicing results in multiple transcript variants.
基因ID	6941

基因名	TCF19
Swiss	Q9Y242
别名	TCF19;SC1;TCF-19

产品验证



Western blot analysis of TCF19 expressed in Mouse thymus,Rat spleen,293T using TCF19 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn