

HDAC11 Rabbit pAb

货号: **AYP16845**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	33kDa/39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,U-87MG,Mouse spinal cord,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 270-347 of human HDA C11 (NP_079103.2).
序列	GLSISPAGIVKRDELVFRMVRGRRVPILMVTSGGYQKRTARIADSILNLFGLGLIGPESPSVSAQNSDTPLLPVAVP

靶点信息

研究背景	This gene encodes a class IV histone deacetylase. The encoded protein is localized to the nucleus and may be involved in regulating the expression of interleukin 10. Alternative splicing results in multiple transcript variants.
基因ID	79885
基因名	HDAC11

Swiss	Q96DB2
别名	HDAC11;HD11

产品验证



Western blot analysis of HDAC11 expressed in Mouse spinal cord,Rat brain,MCF7 using HDAC11 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn