

— ABLYBIO, Help Your Research



SRSF10 Rabbit pAb

货号: **AYP16860**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	20kDa/21kDa/22kDa/31kDa
实测分子量	40KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,22Rv1,K-562,C2C12,Mouse testis,Rat testis
细胞定位	Cytoplasm,Nucleus speckle
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 163-262 of human SRSF10 (NP_473357.1).
------	---

靶点信息

研究背景	This gene product is a member of the serine-arginine (SR) family of proteins, which are involved in constitutive and regulated RNA splicing. Members of this family are characterized by N-terminal RNP1 and RNP2 motifs, which are required for binding to RNA, and multiple C-terminal SR/RS repeats, which are important in mediating association with other cellular proteins. This protein interacts with the oncoprotein TLS, and abrogates the influence of TLS on adenovirus E1A pre-mRNA splicing. This gene has pseudogenes on chromosomes 4, 9, 14, 18, and 20. Alternative splicing results in multiple transcript variants.
基因ID	10772
基因名	SRSF10
Swiss	O75494 (https://www.uniprot.org/uniprotkb/O75494/entry)
别名	SRSF10,FUSIP1,FUSIP2,NSSR,PPP1R149,SFRS13,SFRS13A,SRp38,SRp40,TASR,TASR1,TASR2,SRSF10 Rabbit pAb,40 kDa SR-repressor protein,FUS-interacting serine-arginine-rich protein 1,Splicing factor SRp38,Splicing factor,arginine/serine-rich 13A,TLS-associated protein with Ser-Arg repeats,TLS-associated serine-arginine protein

产品验证



Western blot analysis of SRSF10 expressed in Mouse testis,Rat testis,HeLa using SRSF10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)