

HDLBP Rabbit pAb

货号: **AYP16861**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:200
理论分子量	137kDa/141kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HepG2,Mouse brain,NIH/3T3,Mouse heart,Rat liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1035-1268 of human HDLBP (NP_976221.1).
序列	LERVKELQAEQEDRALRSFKLSVTVDPKYHPKIIGRKGAVITQIRLEHDVNIQFPDKDDGNQPQDQITITGYEKNTEAARDAI LRIVGELEQMVSEDEVPLDHRVHARIIGARGKAIRKIMDEFKVDIRFPQSGAPDPNCVTVTGLPENVEEAIDHILNLEEEYLADV VDSEALQVYMKPPAHEEAKAPSRGFVVRDAPWTASSEKAPDMSSEEFPSFGAQVAPKTLPWGPKR

靶点信息

研究背景	The protein encoded by this gene binds high density lipoprotein (HDL) and may function to regulate excess cholesterol levels in cells. The encoded protein also binds RNA and can induce heterochromatin formation.
------	---

基因ID	3069
基因名	HDLBP
Swiss	Q00341
别名	HDLBP;HBP;PRO2900;VGL;vigilin

产品验证



Western blot analysis of HDLBP expressed in Mouse brain,Rat liver,HeLa using HDLBP Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn