

CD14 Rabbit pAb

货号: **AYP16945**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	40kDa
实测分子量	60KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	RAW264.7
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-330 of human CD14 (NP_001167576.1).
序列	TTPEPCELDDDFRCVCFNFSEFPQPDWSEAFQCVSAVEVEIHAGGLNLEPFLKRVADADADPRQYADTVKALRVRRRLTVGAAQVPAQLLVGALRVLAYSRKELTLEDLKITGTMPPLEATGLALSSLRLRNVSATGRSWLAELQQWLKPKLVLSIAQAHSPAFCSEQVRAFPALTSLDLSDNPGLGERGLMAALCPHKFPAIQNLALRNTGMETPTGVCAALAAAGVQPHSLDLSHNSLRATVNPSPRCMWSSALNSLNSLFAGLEQVPKGLPAKLRVLDLSCNRLNRPQDELPEVDNLTLDGNP

靶点信息

研究背景	The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein.
------	---

基因ID	929
基因名	CD14
Swiss	P08571
别名	CD14

产品验证



Western blot analysis of CD14 expressed in RAW264.7 using CD14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn