

— ABLYBIO, Help Your Research



IRAK3 Rabbit pAb

货号: **AYP17025**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	60kDa/67kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,NCI-H125,Rat liver
细胞定位	cytoplasm,nucleus,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 367-596 of human IRAK 3 (NP_009130.2).
------	--

靶点信息

研究背景	This gene encodes a member of the interleukin-1 receptor-associated kinase protein family. Members of this family are essential components of the Toll/IL-R immune signal transduction pathways. This protein is primarily expressed in monocytes and macrophages and functions as a negative regulator of Toll-like receptor signaling. Mutations in this gene are associated with a susceptibility to asthma. Alternate splicing results in multiple transcript variants.
基因ID	11213
基因名	IRAK3
Swiss	Q9Y616 (https://www.uniprot.org/uniprotkb/Q9Y616/entry)
别名	IRAK3,ASRT5,IRAKM,IRAK3 Rabbit pAb,IL-1 receptor-associated kinase M,Inactive IL-1 receptor-associated kinase 3

产品验证



Western blot analysis of IRAK3 expressed in THP-1, NCI-H125, Rat liver using IRAK3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)