

# HUS1 Rabbit pAb

货号: **AYP17053**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100 <b>IP:</b> 1:50 - 1:200
理论分子量	29kDa/31kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	OVCAR-3,K-562,293T
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human HUS1 ( NP_004498.1).
序列	MKFRKIVDGAACLNHFTRISNMIKLAKTCTLRISPDKLNLFILCDKLANGGVSMWCELEQENFFNEFQMEGVSAENNEIYLE LTSENLSRALKTAQNARALKIKLTNKHFPCLTVSVELLSMSSSRIVTHDIPKVIIPRKLWKDLQEPVVPDPDVSIIYLPVLKTM KSVVEKMKNISNHLVIEANLDGELNLKIETELVCVTTHFKDLGNPPLASESTHEDRNVEHMAEVHIDIRKLLQLAGQQVNP TKALCNIVNNKMHVFDLLHEDVSLQYFIPALS

## 靶点信息

研究背景	The protein encoded by this gene is a component of an evolutionarily conserved, genotoxin-activated checkpoint complex that is involved in the cell cycle arrest in response to DNA damage. This protein forms a heterotrimeric complex with checkpoint proteins RAD9 and RAD1. In response to DNA damage, the trimeric complex interacts with another protein complex consisting of checkpoint protein RAD17 and four small subunits of the replication factor C (RFC), which loads the combined complex onto the chromatin. The DNA damage induced chromatin binding has been shown to depend on the activation of the checkpoint kinase ATM, and is thought to be an early checkpoint signaling event. Alternative splicing results in multiple transcript variants.
基因ID	3364
基因名	HUS1
Swiss	O60921
别名	HUS1;hHUS1

## 产品验证



Western blot analysis of HUS1 expressed in OVCAR-3, K-562, 293T using HUS1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)