

CFP Rabbit pAb

货号: **AYP17057**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:1000 - 1:5000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	51kDa
实测分子量	51KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Human plasma
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 190-469 of human CFP (NP_001138724.1).
序列	CPTHGAWATWGPWTPCSASCHGGPHEPKETRSRKCSAPEPSQKPPGKPCPLAYEQRRCTGLPPCPVAGGWGPWGPV SPCPVTCGLGQTMEQRTC�HPVPQHGGPFAGDATRTHICNTAVPCPVDGEWDSWGEWSPCIRRNMKSISCQEIPGQQ SRGRTCGRKFDGHRCAGQQDIRHCYSIQHCPLKGSWSEWSTWGLCMPPCGPNPTRARQLCTPLLPKYPTVSMVE GQGEKNVTFWGRPLPRCEELQGQKLVEEKRPCLHVPACKDPEEEEL

靶点信息

研究背景	This gene encodes a plasma glycoprotein that positively regulates the alternative complement pathway of the innate immune system. This protein binds to many microbial surfaces and apoptotic cells and stabilizes the C3- and C5-convertase enzyme complexes in a feedback loop that ultimately leads to formation of the membrane attack complex and lysis of the target cell. Mutations in this gene result in two forms of properdin deficiency, which results in high susceptibility to meningococcal infections. Multiple alternatively spliced variants, encoding the same protein, have been identified.
基因ID	5199
基因名	CFP
Swiss	P27918
别名	CFP;BFD;PFC;PFD;PROPERDIN;properdin

产品验证



Western blot analysis of CFP expressed in Human plasma using CFP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn