

— ABLYBIO, Help Your Research



# RAD17 Rabbit pAb

货号: **AYP17076**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:50 - 1:200
理论分子量	57kDa/66kDa/75kDa/77kDa
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,BT-474,K-562,22Rv1,Raji
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 391-670 of human RAD 17 (NP_002864.1).
------	----------------------------------------------------------------------------------------------------------------------

## 靶点信息

研究背景	The protein encoded by this gene is highly similar to the gene product of <i>Schizosaccharomyces pombe</i> rad17, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This protein shares strong similarity with DNA replication factor C (RFC), and can form a complex with RFCs. This protein binds to chromatin prior to DNA damage and is phosphorylated by the checkpoint kinase ATR following damage. This protein recruits the RAD1-RAD9-HUS1 checkpoint protein complex onto chromatin after DNA damage, which may be required for its phosphorylation. The phosphorylation of this protein is required for the DNA-damage-induced cell cycle G2 arrest, and is thought to be a critical early event during checkpoint signaling in DNA-damaged cells. Multiple alternatively spliced transcript variants of this gene, which encode four distinct protein isoforms, have been reported. Two pseudogenes, located on chromosomes 7 and 13, have been identified.
基因ID	5884
基因名	RAD17
Swiss	O75943 ( <a href="https://www.uniprot.org/uniprotkb/O75943/entry">https://www.uniprot.org/uniprotkb/O75943/entry</a> )
别名	RAD17,CCYC,HRAD17,R24L,RAD17SP,RAD24,RAD17 Rabbit pAb,RF-C/activator 1 homolog

## 产品验证



Western blot analysis of RAD17 expressed in SW480,BT-474,K-562 using RAD17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)