

— ABLYBIO, Help Your Research



NMNAT1 Rabbit pAb

货号: **AYP17126**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	31kDa
实测分子量	29KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human NMNAT 1 (NP_073624.2).
------	---

靶点信息

研究背景	This gene encodes an enzyme which catalyzes a key step in the biosynthesis of nicotinamide adenine dinucleotide (NAD). The encoded enzyme is one of several nicotinamide nucleotide adenylyltransferases, and is specifically localized to the cell nucleus. Activity of this protein leads to the activation of a nuclear deacetylase that functions in the protection of damaged neurons. Mutations in this gene have been associated with Leber congenital amaurosis 9. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are located on chromosomes 1, 3, 4, 14, and 15.
基因ID	64802
基因名	NMNAT1
Swiss	Q9HAN9 (https://www.uniprot.org/uniprotkb/Q9HAN9/entry)
别名	NMNAT1,LCA9,NMNAT,PNAT1,NMNAT1 Rabbit pAb,Nicotinamide-nucleotide adenylyltransferase 1,Nicotinate-nucleotide adenylyltransferase 1

产品验证



Western blot analysis of NMNAT1 expressed in 293T using NMNAT1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)