

TPK1 Rabbit pAb

货号: **AYP17153**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	13kDa/27kDa
实测分子量	27KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Rat kidney
细胞定位	cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-243 of human TPK1 (NP_071890.2).
序列	MEHAFTPLEPLLSTGNLKYCLVILNQPLDNYFRHLWNLKALLRACADGGANRLYDITEGERESFLPEFINGDFDSIRPEVREYY ATKGCELISTPDQDHTDFTKCLKMLQKKIEEKDLKVDVIVTLGGLAGRFQIMASVNTLFQATHITPFPIIIQEESELIYLLQPG KHRLHVDTGMEGDWCGLIPVGQPCMQVTTTGLKWNLNDVLAFTLVSTSNYDGSVVTVETDHPLLWTMAIKS

靶点信息

研究背景	The protein encoded by this gene functions as a homodimer and catalyzes the conversion of thiamine to thiamine pyrophosphate, a cofactor for some enzymes of the glycolytic and energy production pathways. Defects in this gene are a cause of thiamine metabolism dysfunction syndrome-5.
基因ID	27010

基因名	TPK1
Swiss	Q9H3S4
别名	TPK1;HTPK1;PP20;THMD5

产品验证



Western blot analysis of TPK1 expressed in Mouse kidney,Rat kidney using TPK1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn