

BLOC1S6 Rabbit pAb

货号: AYP17156

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	8kDa/19kDa
实测分子量	20kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HepG2,DU145,K-562,Rat testis
细胞定位	Cytoplasm,Endomembrane system,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-170 of human BLOC1 S6 (NP_036520.1).
序列	MSVPGPSSPDGALTRPPYCLEAGEPTPLSDTSPDEGLIEDLTIEDKAVEQLAEGLLSHYLPDLQRSKQALQELTQNQVLLDTLEQEISKFKCHECHSMLDINALFAEAKHYHAKLVNIRKEMMLHEKTSKLLKRALKLQQKRQKEELEREQQREKEFEREKQLTARPAK

靶点信息

研究背景	The protein encoded by this gene may play a role in intracellular vesicle trafficking. It interacts with Syntaxin 13 which mediates intracellular membrane fusion. Mutations in this gene cause symptoms associated with Hermansky-Pudlak syndrome-9. Alternative splicing results in multiple transcript variants. A pseudogene related to this gene is located on the X chromosome.
基因ID	26258

基因名	BLOC1S6
Swiss	Q9UL45
别名	BLOC1S6;BLOS6;HPS9;PA;PALLID;PLDN

产品验证



Western blot analysis of BLOC1S6 expressed in Rat testis,U-87MG,HepG2 using BLOC1S6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn