

— ABLYBIO, Help Your Research



LARP1 Rabbit pAb

货号: **AYP17169**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	116kDa/123kDa
实测分子量	163kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 106-205 of human LARP 1 (Q6PKG0).
------	---

靶点信息

研究背景	RNA-binding protein that regulates the translation of specific target mRNA species downstream of the mTORC1 complex, in function of growth signals and nutrient availability. Interacts on the one hand with the 3' poly-A tails that are present in all mRNA molecules, and on the other hand with the 7-methylguanosine cap structure of mRNAs containing a 5' terminal oligopyrimidine (5'TOP motif, which is present in mRNAs encoding ribosomal proteins and several components of the translation machinery. The interaction with the 5' end of mRNAs containing a 5'TOP motif leads to translational repression by preventing the binding of EIF4G1. When mTORC1 is activated, LARP1 is phosphorylated and dissociates from the 5' untranslated region (UTR of mRNA. Does not prevent binding of EIF4G1 to mRNAs that lack a 5'TOP motif. Interacts with the free 40S ribosome subunit and with ribosomes, both monosomes and polysomes. Under normal nutrient availability, interacts primarily with the 3' untranslated region (UTR of mRNAs encoding ribosomal proteins and increases protein synthesis. Associates with actively translating ribosomes and stimulates translation of mRNAs containing a 5'TOP motif, thereby regulating protein synthesis, and as a consequence, cell growth and proliferation. Stabilizes mRNAs species with a 5'TOP motif, which is required to prevent apoptosis.
基因ID	23367
基因名	LARP1
Swiss	Q6PKG0 (https://www.uniprot.org/uniprotkb/Q6PKG0/entry)
别名	LARP1,LARP,LARP1 Rabbit pAb,La ribonucleoprotein domain family member 1,KIAA0731

产品验证



Western blot analysis of LARP1 expressed in HeLa, MCF7 using LARP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)