

FCRL3 Rabbit pAb

货号: **AYP17182**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	21kDa/22kDa/69kDa/77kDa/80kDa/81kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,B cells,293T,U-937,Mouse kidney,Rat kidney
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 600-720 of human FCRL 3 (NP_443171.2).
序列	RRKPGGLSATGTSSHSPSECQEPSSSRPSRIDPQEPHSHKPLAPMELEPMYSNVNPGDSNPIYSQIWSIQHTKENSANCPM MHQEHEELTVLYSELKKTHPDSSAGEASSRGRAHEEDDEE

靶点信息

研究背景	This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein contains immunoreceptor-tyrosine activation motifs and immunoreceptor-tyrosine inhibitory motifs in its cytoplasmic domain and may play a role in regulation of the immune system. Mutations in this gene have been associated with rheumatoid arthritis, autoimmune thyroid disease, and systemic lupus erythematosus. Alternative splicing results in multiple transcript variants.
------	---

基因ID	115352
基因名	FCRL3
Swiss	Q96P31
别名	FCRL3;CD307c;FCRH3;IFGP3;IRTA3;SPAP2

产品验证



Western blot analysis of FCRL3 expressed in Mouse kidney,Rat kidney,Raji using FCRL3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn