

— ABLYBIO, Help Your Research



# FCRL3 Rabbit pAb

货号: **AYP17182**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB IHC   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IHC:</b> 1:100 - 1:200   |
| 理论分子量 | 21kDa/22kDa/69kDa/77kDa/80kDa/81kDa  |
| 实测分子量 | 105kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | Raji,B cells,293T,U-937,Mouse kidney,Rat kidney  |
| 细胞定位  | Cell membrane,Single-pass type I membrane protein  |
| 纯化    | Affinity purification  |


## 抗原信息

|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 600-720 of human FCRL 3 (NP_443171.2). |
|------|--|

## 靶点信息

|       |   |
|-------|---|
| 研究背景  | This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein contains immunoreceptor-tyrosine activation motifs and immunoreceptor-tyrosine inhibitory motifs in its cytoplasmic domain and may play a role in regulation of the immune system. Mutations in this gene have been associated with rheumatoid arthritis, autoimmune thyroid disease, and systemic lupus erythematosus. Alternative splicing results in multiple transcript variants. |
| 基因ID  | 115352  |
| 基因名   | FCRL3   |
| Swiss | Q96P31 ( <a href="https://www.uniprot.org/uniprotkb/Q96P31/entry">https://www.uniprot.org/uniprotkb/Q96P31/entry</a> )  |
| 别名    | FCRL3,CD307c,FCRH3,IFGP3,IRTA3,SPAP2,FCRL3 Rabbit pAb,Fc receptor homolog 3,IFGP family protein 3,Immune receptor translocation-associated protein 3,MAIA,SH2 domain-containing phosphatase anchor protein 2  |

## 产品验证



Western blot analysis of FCRL3 expressed in Mouse kidney,Rat kidney,Raji using FCRL3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)