

GOSR2 Rabbit pAb

货号: AYP17205

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	22kDa/24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A-549, HT-29, MCF7, HT-1080, HepG2, K-562
细胞定位	Golgi apparatus, Golgi apparatus membrane, Single-pass type IV membrane protein, cis-Golgi network membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human GOSR2 (NP_004278.2).
序列	MDPLFQQTHKQVHEIQSCMGRLETADKQSVHIVENEIQASIDQIFSRLERLEILSSKEPPNKRQNARLRVDQLKYDVQHLQ TALRNFQHRRHAREQERQREELLSRTFTTNDSDTTIPMDESLLQFNSSLQKVHNGMDDLILDGHNILDGLRTQRLTLKGT QKKILDIANMLGLSNTVMRLIEKRAFQDK

靶点信息

研究背景	This gene encodes a trafficking membrane protein which transports proteins among the medial- and trans-Golgi compartments. Due to its chromosomal location and trafficking function, this gene may be involved in familial essential hypertension.
基因ID	9570

基因名	GOSR2
Swiss	O14653
别名	GOSR2;Bos1;EPM6;GS27

产品验证



Western blot analysis of GOSR2 expressed in A-549, HT-29, MCF7 using GOSR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn