

— ABLYBIO, Help Your Research



# EIF1AY Rabbit pAb

货号: **AYP17212**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:200 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	16kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,MCF7,Raji,Jurkat,U-251MG,Mouse brain,Rat brain
细胞定位	cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-144 of human EIF1AY (NP_004672.2).
------	--

## 靶点信息

研究背景	This gene is located on the non-recombining region of the Y chromosome. It encodes a protein related to eukaryotic translation initiation factor 1A (EIF1A), which may function in stabilizing the binding of the initiator Met-tRNA to 40S ribosomal subunits. Alternative splicing results in multiple transcript variants.
基因ID	9086
基因名	EIF1AY
Swiss	O14602 ( <a href="https://www.uniprot.org/uniprotkb/O14602/entry">https://www.uniprot.org/uniprotkb/O14602/entry</a> )
别名	EIF1AY,eIF-4C,Y-linked,EIF1AY Rabbit pAb,Eukaryotic translation initiation factor 4C

## 产品验证



Western blot analysis of EIF1AY expressed in Mouse brain,Rat brain,NIH/3T3 using EIF1AY Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)