

— ABLYBIO, Help Your Research



# RBM10 Rabbit pAb

货号: **AYP17221**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	94kDa/103kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat, MCF7, LO2, 293T
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human RBM10 (NP_005667.2).
------	---

## 靶点信息

研究背景	This gene encodes a nuclear protein that belongs to a family proteins that contain an RNA-binding motif. The encoded protein associates with hnRNP proteins and may be involved in regulating alternative splicing. Defects in this gene are the cause of the X-linked recessive disorder, TARP syndrome. Alternate splicing results in multiple transcript variants.
基因ID	8241
基因名	RBM10
Swiss	P98175 ( <a href="https://www.uniprot.org/uniprotkb/P98175/entry">https://www.uniprot.org/uniprotkb/P98175/entry</a> )
别名	RBM10,DXS8237E,GPATC9,GPATCH9,S1-1,TARPS,ZRANB5,RBM10 Rabbit pAb,G patch domain-containing protein 9,RNA-binding motif protein 10,RNA-binding protein S1-1,KIAA0122

## 产品验证



Western blot analysis of RBM10 expressed in Jurkat, MCF7, LO2 using RBM10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)