

— ABLYBIO, Help Your Research



PRKCSH Rabbit pAb

货号: **AYP17245**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	59kDa
实测分子量	80KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,A-549,Mouse lung,Rat brain
细胞定位	Endoplasmic reticulum
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 130-275 of human PRKCSH (P14314).
------	---

靶点信息

研究背景	This gene encodes the beta-subunit of glucosidase II, an N-linked glycan-processing enzyme in the endoplasmic reticulum. The encoded protein is an acidic phosphoprotein known to be a substrate for protein kinase C. Mutations in this gene have been associated with the autosomal dominant polycystic liver disease. Alternative splicing results in multiple transcript variants.
基因ID	5589
基因名	PRKCSH
Swiss	P14314 (https://www.uniprot.org/uniprotkb/P14314/entry)
别名	PRKCSH,AGE-R2,G19P1,GIIB,PCLD,PCLD1,PKCSH,PLD1,VASAP-60,PRKCSH Rabbit pAb,80K-H protein,Glucosidase II subunit beta,Protein kinase C substrate 60.1 kDa protein heavy chain

产品验证



Western blot analysis of PRKCSH expressed in Mouse lung,Rat brain,Jurkat using PRKCSH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)