

FUT6 Rabbit pAb

货号: **AYP17262**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	41kDa/42kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480
细胞定位	Golgi apparatus,Golgi stack membrane,Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 170-359 of human FUT6 (NP_000141.1).
序列	WLEPWGQPAHPLNLSAKTELVAWAVSNWGPNSARVRYYSQSLQAHLKVDVYGRSHKPLPQGTMMETLSRYKFYLAFE NSLHPDYITEKLWRNALEAWAVPVVLGSPRSNYERFLPPDAFIHVDDFQSPKDLARYLQELDKDHARYLSYFRWRETLRPR SFSWALAFCKACWKLQEESRYQTRGIAAWFT

靶点信息

研究背景	The protein encoded by this gene is a Golgi stack membrane protein that is involved in the creation of sialyl-Lewis X, an E-selectin ligand. Mutations in this gene are a cause of fucosyltransferase-6 deficiency. Two transcript variants encoding the same protein have been found for this gene.
基因ID	2528

基因名	FUT6
Swiss	P51993
别名	FUT6;FCT3A;FT1A;Fuc-TVI;FucT-VI

产品验证



Western blot analysis of FUT6 expressed in SW480 using FUT6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn