

EIF2S3 Rabbit pAb

货号: **AYP17267**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	51kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	cytoplasm, cytosol, eukaryotic translation initiation factor 2 complex, extracellular exosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 173-472 of human EIF2 S3 (NP_001406.1).
序列	HLAAIEIMKCLKHILQNKIDLVKESQAKEQYEQILAFVQGTVAEGAPIIPISAQLKYNIEVVCEYIVKKIPVPPRDFTEPRLIVIRSF FDVKNKPGCEVDDLKGGVAGGSILKGVKVGQEIEVRPGIVSKDSEGKLMCKPIFSKIVSLFAEHNDLQYAAPGGLIGVGT KI DPTLCRADRMVGVQVLGAVGALPEIFTELEISYFLLRRLGVRTEGDKKAAKVQKLSKNEVLMVNIGSLSTGGRVSAVKADL GKIVLTNPVCTEVGEKIALSRRVEKHWRLIGWQIRRGVTIKPTVDDD

靶点信息

研究背景	The protein encoded by this gene is the largest subunit of a heterotrimeric GTP-binding protein involved in the recruitment of methionyl-tRNA(i) to the 40 S ribosomal subunit.
基因ID	1968

基因名	EIF2S3
Swiss	P41091
别名	EIF2S3;EIF2;EIF2G;EIF2gamma;MRXSBRK;eIF-2gA

产品验证



Western blot analysis of EIF2S3 expressed in Rat brain using EIF2S3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn