

— ABLYBIO, Help Your Research



ARHGAP1 Rabbit pAb

货号: **AYP17288**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:50 - 1:100
理论分子量	50kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,HeLa,HepG2,U-87MG,PC-12,Mouse lung
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 210-439 of human ARHGAP1 (NP_004299.1).
------	---

靶点信息

研究背景	This gene encodes a member of a large family of proteins that activate Rho-type guanosine triphosphate (GTP) metabolizing enzymes. The encoded protein contains a SRC homology 3 domain and interacts with Bcl-2-associated protein family members.
基因ID	392
基因名	ARHGAP1
Swiss	Q07960 (https://www.uniprot.org/uniprotkb/Q07960/entry)
别名	ARHGAP1,CDC42GAP,RHOGAP,RHOGAP1,p50rhoGAP,ARHGAP1 Rabbit pAb,CDC42 GTPase-activating protein,GTPase-activating protein rhoGAP,Rho-related small GTPase protein activator,Rho-type GTPase-activating protein 1,p50-RhoGAP

产品验证



Western blot analysis of ARHGAP1 expressed in Mouse lung, BxPC-3, HeLa using ARHGAP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)