

THOC2 Rabbit pAb

货号: **AYP17305**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	183kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse liver
细胞定位	nuclear speck,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human THOC2 (NP_001075019.1).
序列	MAAAVVVPAEWIKNWEKSGRGEFLHLCRILSENKSHDSSTYRDFQQALYELSYHVIKGNLKHQASNVLSDISEFREDM PSILADVFCILDITNCLLEEKSKRDYFTQLVLAACYLVSDTVLKERLDPETLESGLIKQSQQFNQKSVKIKTKLFYKQKFNLL REENEGYAKLIAELGQDLSGSITSDLILENIKSLIGCFNLDPNRVLDVILEVFECRPEHDDFFISLLESYMSMCEPQTLCHILGF KFKFYQEPNGETPSSLYRVA

靶点信息

研究背景	The TREX multiprotein complex binds specifically to spliced mRNAs to facilitate mRNA export. The protein encoded by this gene is a member of the THO complex, a subset of the TREX complex. The encoded protein interacts with the THOC1 protein.
基因ID	57187

基因名	THOC2
Swiss	Q8NI27
别名	THO2;MRX12;MRX35;CXorf3;hTREX120;dj506G2.1;THOC2

产品验证



Western blot analysis of THOC2 expressed in HeLa, Mouse liver using THOC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn