

SLITRK6 Rabbit pAb

货号: **AYP17306**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	95kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse brain,Rat brain
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 621-816 of human SLITRK6 (NP_115605.2).
序列	IVFCAAGIVVLVLRHRRRYYKKKQVDEQMRDNSPVHLQYSMYGHKTHHTTTERPSASLYEQHMVSPMVHVYRSPSFGPKHLEEEEEERNEKEGSDAKHLQRSLLLEQENHSPLTGSNMKYKTTNQSTEFLSFQDASSLYRNILEKERELQQLGITEYLRKNIAQLQPDMEAHYPGAHEELKLMETLMYSRPRKVLVEQT

靶点信息

研究背景	This gene encodes a member of the SLITRK protein family. Members of this family are integral membrane proteins that are characterized by two N-terminal leucine-rich repeat (LRR) domains and a C-terminal region that shares homology with trk neurotrophin receptors. This protein functions as a regulator of neurite outgrowth required for normal hearing and vision. Mutations in this gene are a cause of myopia and deafness.
------	---

基因ID	84189
基因名	SLITRK6
Swiss	Q9H5Y7
别名	SLITRK6;DFNMYP

产品验证



Western blot analysis of SLITRK6 expressed in 293T, Mouse brain, Rat brain using SLITRK6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn