

C14orf166 Rabbit pAb

货号: **AYP17313**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 28kDa |
| 实测分子量 | 25kDa/28kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Jurkat,293T,NIH/3T3 |
| 细胞定位 | Cytoplasm,Nucleus,centrosome,cytoskeleton,cytosol,microtubule organizing center,perinuclear region |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-244 of human C14orf166 (NP_057123.1). |
| 序列 | MFRRKLTALDYHNPAGFNCKDETEFRNFIVWLEDQKIRHYKIEDRGNLRNIHSSDWPKEFEKYL RDVNC PFKIQDRQE AIDWLLGLAVRLEYGDNAEKYKDLVDPNSKTADNATKNAEPLINLDVNNPDFKAGVMALANLLQIRHDDYL VMLKAIRILVQE RLTQDAVAKANQTK EGLPVALDKHILGFDTGDAVLN EAAQILRLLHIEELRELQTKINEAIVAVQAI IADPKTDHRLGKVGR |

靶点信息

| | |
|------|--|
| 研究背景 | RNA-binding protein involved in modulation of mRNA transcription by Polymerase II. Component of the tRNA-splicing ligase complex and is required for tRNA ligation. May be required for RNA transport. |
| 基因ID | 51637 |

| | |
|-------|---|
| 基因名 | C14orf166 |
| Swiss | Q9Y224 |
| 别名 | C14orf166;CGI-99;CGI99;CLE;CLE7;LCRP369;RLLM1;hCLE1 |

产品验证



Western blot analysis of C14orf166 expressed in Jurkat,293T,NIH/3T3 using C14orf166 Rabbit pAb at 1 :1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn