

# BZW1 Rabbit pAb

货号: **AYP17320**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	40kDa/48kDa/51kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,LO2,BT-474,Mouse liver,Mouse kidney
细胞定位	cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-290 of human BZW1 (NP_001193996.1).
序列	MNNQKQKPTLSGQRFKTRKRDEKERFDPTQFQDCIIQGLTETGTDLAVAKFLDASGAKLDYRRYAETLFDILVAGGMLAPGGTLADDDMMRTDVCVFAAQEDLETMQAFAQVFNKLIRRYKYLEKGFEDVKKLLFLKGFSESERNKLAMLTGVLLANGTLNASILNSLYNENLVKEGVSAAFVVKLFSWINEKDINAVAASLRKVSMNRLMELFPANKQSVHEFTKYFTEAGLKELSEYVRNQQTIGARKELQKELQE QMSRGDPFKDIIYVKEEMKNNIPEP

## 靶点信息

研究背景	Enhances histone H4 gene transcription but does not seem to bind DNA directly.
基因ID	9689
基因名	BZW1

<b>Swiss</b>	Q7L1Q6
别名	BZW1;BZAP45;Nbla10236

## 产品验证



Western blot analysis of BZW1 expressed in Mouse liver, HeLa, LO2 using BZW1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)