

RBM17 Rabbit pAb

货号: **AYP17345**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:200 - 1:2000 IHC: 1:50 - 1:200
理论分子量	44kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,A-549,HeLa,DU145,Rat thymus
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human RBM17 (NP_116294.1).
序列	MSLYDDLGVETSDSKTEGWSKNFKLLQSQLQVKKAAALTQAKSQRTKQSTVLAPVIDLKRGGSSDDRQIVDTPPHVAAGL KDPVPSGFSAGEVLIPLADEYDPMFPNDYEKVVKRQREERQQRQLERQKEIEEREKRRKDRHEASGFARRPDPDSDEDE DYERERRKRSMGGAAIAPPTSLVEKDKELPRDFPYEEDSRPRSQSSKAAIPPPVYEEQDRPRSPTGPSNSFLANMGGTVAH

靶点信息

研究背景	This gene encodes an RNA binding protein. The encoded protein is part of the spliceosome complex and functions in the second catalytic step of mRNA splicing. Alternatively spliced transcript variants have been described. Related pseudogenes exist on chromosomes 9 and 15.
基因ID	84991

基因名	RBM17
Swiss	Q96I25
别名	RBM17;SPF45

产品验证



Western blot analysis of RBM17 expressed in Rat thymus, LO2, A-549 using RBM17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn