

— ABLYBIO, Help Your Research



MC1 Receptor Rabbit pAb

货号: **AYP17394**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	34kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	M21,MCF7,SW620,22Rv1,Raji,Mouse testis
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant protein of human MC1 Receptor
------	---

靶点信息

研究背景	This intronless gene encodes the receptor protein for melanocyte-stimulating hormone (MSH). The encoded protein, a seven pass transmembrane G protein coupled receptor, controls melanogenesis. Two types of melanin exist: red pheomelanin and black eumelanin. Gene mutations that lead to a loss in function are associated with increased pheomelanin production, which leads to lighter skin and hair color. Eumelanin is photoprotective but pheomelanin may contribute to UV-induced skin damage by generating free radicals upon UV radiation. Binding of MSH to its receptor activates the receptor and stimulates eumelanin synthesis. This receptor is a major determining factor in sun sensitivity and is a genetic risk factor for melanoma and non-melanoma skin cancer. Over 30 variant alleles have been identified which correlate with skin and hair color, providing evidence that this gene is an important component in determining normal human pigment variation.
基因ID	4157
基因名	MC1R
Swiss	Q01726 (https://www.uniprot.org/uniprotkb/Q01726/entry)
别名	MC1R,CMM5,MSH-R,SHEP2,MC1 Receptor Rabbit pAb,Melanocortin receptor 1,MSHR

产品验证



Western blot analysis of MC1 Receptor expressed in Mouse testis, M21, MCF7 using MC1 Receptor Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)