

GFRA3 Rabbit pAb

货号: **AYP17413**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	40kDa/44kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,BxPC-3,HepG2
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 32-240 of human GFRA3 (NP_001487.2).
序列	DPLPTESRLMNSCLQARRKCQADPTCSAAYHHLDSCTSSISTPLPSEEPSVPADCLEAAQQLRNSSLIGCMCHRRMKNQV ACLDIYWTVHRARSLGNYELDVSPYEDTVTSKPWKMNLSKLNMLKPDSDLCLKFAMLCTLNDKCDRLRKAYGEACSGPH CQRHVCLRQLLTFEKAEPHAQGLLLCPCAPNDRGCGERRRNTIAPNCA

靶点信息

研究背景	The protein encoded by this gene is a glycosylphosphatidylinositol(GPI)-linked cell surface receptor and a member of the GDNF receptor family. It forms a signaling receptor complex with RET tyrosine kinase receptor and binds the ligand, artemin.
基因ID	2676

基因名	GFRA3
Swiss	O60609
别名	GFRA3;GDNFR3

产品验证



Western blot analysis of GFRA3 expressed in A-549, BxPC-3, HepG2 using GFRA3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn