

— ABLYBIO, Help Your Research



Gelsolin Rabbit pAb

货号: **AYP17415**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	80kDa/81kDa/85kDa
实测分子量	109kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,SW480
细胞定位	Cytoplasm,Secreted,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 543-782 of human Gelsolin (NP_000168.1).
------	--

靶点信息

研究背景	The protein encoded by this gene binds to the 'plus' ends of actin monomers and filaments to prevent monomer exchange. The encoded calcium-regulated protein functions in both assembly and disassembly of actin filaments. Defects in this gene are a cause of familial amyloidosis Finnish type (FAF). Multiple transcript variants encoding several different isoforms have been found for this gene.
基因ID	2934
基因名	GSN
Swiss	P06396 (https://www.uniprot.org/uniprotkb/P06396/entry)
别名	GSN,ADF,AGEL,gelsolin,Gelsolin Rabbit pAb,Actin-depolymerizing factor,Brevin

产品验证



Western blot analysis of Gelsolin expressed in U-251MG,SW480 using Gelsolin Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)