

# CXCR2 Rabbit pAb

货号: AYP17430

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	40kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HT-29
细胞定位	Cell membrane, Multi-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human CXCR2 (P25025).
序列	MEDFNMESDSFEDFWKGEDLSNYSYSSTLPPFLDAAAPCEPESLEINKYFVVIYALVFLLSLLGNSLVMLVILYSRVGRSVT DVYLLNLALADLLFALT

## 靶点信息

研究背景	The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. Alternatively spliced variants, encoding the same protein, have been identified.
基因ID	3579
基因名	CXCR2
Swiss	P25025
别名	CXCR2;CD182;CDw128b;CMKAR2;IL8R2;IL8RA;IL8RB

## 产品验证



Western blot analysis of CXCR2 expressed in HT-29 using CXCR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)