

DNAJC7 Rabbit pAb

货号: **AYP17453**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	50kDa/56kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,293T,K-562
细胞定位	Cytoplasm,Nucleus,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 230-380 of human DNAJ C7 (NP_003306.3).
序列	VQFFVQALRMAPDHEKACIACRNAKALKAKKEDGNKAFKEGNYKLAYELYTEALGIDPNNIKTNAKLYCNRGTVNSKLRKLD DDAIEDCTNAVKLDDTYIKAYLRRRAQCYMDTEQYEEAVRDYKVVYQTEKTEKHKQLLKNAQLELKKSKRK

靶点信息

研究背景	This gene encodes a member of the DNAJ heat shock protein 40 family of proteins that is characterized by two N-terminal tetratricopeptide repeat domains and a C-terminal DNAJ domain. This protein binds the chaperone proteins heat shock proteins 70 and 90 in an ATP-dependent manner and may function as a co-chaperone. Pseudogenes of this gene are found on chromosomes 1 and 6. Alternate splicing results in multiple transcript variants.
基因ID	7266

基因名	DNAJC7
Swiss	Q99615
别名	DNAJC7;DJ11;DJC7;TPR2;TTC2

产品验证



Western blot analysis of DNAJC7 expressed in A-431,293T,K-562 using DNAJC7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn