

IRAK2 Rabbit pAb

货号: **AYP17457**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	69kDa
实测分子量	78kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	cytoplasm,cytosol,nucleus,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human IRAK2 (NP_001561.3).
序列	MACYIYQLPSWVLDLDCRNMDALSEWDWMEFASYVITDLTQLRKIKSMERVQGV SITRELLWWWGMRQATVQQLVDLL CRLELYRAAQIILNWKPAPEIRCPIPAFPDSVKPEKPLAASVRKAEDEQEEGQPVRMATFPGPGSSPARAHQPAFLQPPEED APHSLSLDLPTSSDSKDFSTSIPKQEKLLSLAGDSLFWSEADV VQATDDFNQNRKISQGT FADVVRGHRHGKPFVFKCLR ETACSSPGSIERFFQAE LQICLRCHPNVL

靶点信息

研究背景	IRAK2 encodes the interleukin-1 receptor-associated kinase 2, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. IRAK2 is reported to participate in the IL1-induced upregulation of NF-kappaB.
基因ID	3656

基因名	IRAK2
Swiss	O43187
别名	IRAK2;IRAK-2

产品验证



Western blot analysis of IRAK2 expressed in Mouse liver using IRAK2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn