

— ABLYBIO, Help Your Research



# HNMT Rabbit pAb

货号: **AYP17510**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100 <b>IP:</b> 1:50 - 1:200
理论分子量	6kDa/14kDa/33kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	22RV1,NCI-H460,HT-29,HepG2,HeLa,Mouse liver,Mouse brain
细胞定位	Cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-292 of human HNMT (NP_008826.1).
------	--

## 靶点信息

研究背景	In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different proteins have been found for this gene.
基因ID	3176
基因名	HNMT
Swiss	P50135 ( <a href="https://www.uniprot.org/uniprotkb/P50135/entry">https://www.uniprot.org/uniprotkb/P50135/entry</a> )
别名	HNMT,HMT,HNMT-S1,HNMT-S2,MRT51,HNMT Rabbit pAb

## 产品验证



Western blot analysis of HNMT expressed in Mouse liver,22RV1,NCI-H460 using HNMT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))