

PDCD4 Rabbit pAb

货号: **AYP17532**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100 IP: 1:20 - 1:50
理论分子量	50kDa/51kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human PDCD4 (NP_055271.2).
序列	MDVENEQILNVNPADPDNLSDSLFSGDEENAGTEEIKNEINGNWISASSINEARINAKAKRRLRKNSRDSGRGDSVSDS GSDALRSGTLVPTSPKGRLLDRRSRSGKGRGLPKKGGAGGKGVWGTGPGQVYDVEEVDVKDPNYDDDDQENCVYETVVL PLDERAFEKTLTPIIQEYFEHGDTNEVAEMLRDLNLGEMKSGVPLAVSLALEGKASHREMTSKLLSDLCGTVMSTTDVEK SFDKLLKDLPELALDTPRAPQL

靶点信息

研究背景	This gene is a tumor suppressor and encodes a protein that binds to the eukaryotic translation initiation factor 4A1 and inhibits its function by preventing RNA binding. Alternative splicing results in multiple transcript variants.
------	---

基因ID	27250
基因名	PDCD4
Swiss	Q53EL6
别名	PDCD4;H731

产品验证



Western blot analysis of PDCD4 expressed in HeLa, MCF7 using PDCD4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn