

— ABLYBIO, Help Your Research



# KCNQ1 Rabbit pAb

货号: **AYP17559**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	61kDa/74kDa
实测分子量	70KDa/75KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse pancreas,Rat lung,Rat kidney,Rat heart
细胞定位	Cell membrane,Cytoplasmic vesicle membrane,Multi-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-549 of human KCNQ1 (NP_861463.1).
------	---

## 靶点信息

研究背景	This gene encodes a voltage-gated potassium channel required for repolarization phase of the cardiac action potential. This protein can form heteromultimers with two other potassium channel proteins, KCNE1 and KCNE3. Mutations in this gene are associated with hereditary long QT syndrome 1 (also known as Romano-Ward syndrome), Jervell and Lange-Nielsen syndrome, and familial atrial fibrillation. This gene exhibits tissue-specific imprinting, with preferential expression from the maternal allele in some tissues, and biallelic expression in others. This gene is located in a region of chromosome 11 amongst other imprinted genes that are associated with Beckwith-Wiedemann syndrome (BWS), and itself has been shown to be disrupted by chromosomal rearrangements in patients with BWS. Alternatively spliced transcript variants have been found for this gene.
基因ID	3784
基因名	KCNQ1
Swiss	P51787 ( <a href="https://www.uniprot.org/uniprotkb/P51787/entry">https://www.uniprot.org/uniprotkb/P51787/entry</a> )
别名	KCNQ1,ATFB1,ATFB3,JLNS1,KCNA8,KCNA9,KVLQT1,Kv1.9,Kv7.1,LQT,LQT1,RWS,SQT2,WRS,KCNQ1 Rabbit pAb,IKs producing slow voltage-gated potassium channel subunit alpha KvLQT1,KQT-like 1,Voltage-gated potassium channel subunit Kv7.1

## 产品验证



Western blot analysis of KCNQ1 expressed in Mouse pancreas, Rat lung, 293T using KCNQ1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)