

HCLS1 Rabbit pAb

货号: **AYP17563**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	24kDa/54kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,Jurkat
细胞定位	Cytoplasm,Membrane,Mitochondrion,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human HCLS1 (NP_005326.2).
序列	MWKSVVGHVSVSVETQGDDWDTDPDFVNDISEKEQRWGAKTIEGSGRTEHINIHLRNKVSEEHVLRKKEMESGPK ASHGYGGRFGVERDRMDKSAVGHEHYVAEVEKHSSQTDAAKGFGGKYGVERDRADKSAVGFDYKGEVEKHTSQKDYSR GFGGRYGVKDKWDKAALGYDYKGETEKHESQRDYAKGFGGQYGIQKDRVDKSAVGFNEMEAPTTAYKKTTPIEAASS GTRGLKAKFESMAEEKRKREEEKAQQVARRQKERKAVTKRSPEAPQ

靶点信息

研究背景	HCLS1 (hematopoietic cell-specific Lyn substrate 1) is a protein-coding gene. Diseases associated with HCLS1 include chronic lymphocytic leukemia. GO annotations related to this gene include protein kinase binding and protein complex binding. An important paralog of this gene is DBN1.
------	---

基因ID	3059
基因名	HCLS1
Swiss	P14317
别名	HCLS1;CTTNL;HS1;lckBP1;p75

产品验证



Western blot analysis of HCLS1 expressed in BT-474, Jurkat using HCLS1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn