

Egr4 Rabbit pAb

货号: **AYP17592**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	61kDa
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain, Mouse kidney
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-367 of mouse Egr4 (NP_065621.1).
序列	MLHLSDFSGSDALLSKPTEGCAHASPELPRLPARDAPSAAAYPGGDFLSWALSSCGAGGDLTDSCFLEGPAPTPPSGLSY SGSFFIQAVPEHPHDPEALFNLMGILGLAPFPGPEAAASRPLDVFPAGPDALLPDLYSPDLSSAAFPEAFWEAAPSAGA PSQCLFEPQLSPDVKPLRAPPASPALDAAASAFKGPYAPWELLSAGVPGNCGSQGSFQTTPPEARFSAVGKVEDLLSIS CPAELPGPAARLYQAGAYDTFSLAPGDLGEGTEGLPALLTPPGGEGGSGGEGGEFLAAPPAQLSPLGLRGAATADFSKPL VADLPGGSGVAAPSSPAASFPAKARRKRRGGKCSARCF CPRP

靶点信息

研究背景	Transcriptional regulator. Recognizes and binds to the DNA sequence 5'-GCGGGGGCG-3' (GSG). Activates the transcription of target genes whose products are required for mitogenesis and differentiation (By similarity).
------	---

基因ID	13656
基因名	
Swiss	Q05215
别名	EGR4;NGFI-C;NGFIC;PAT133;NGF1-C;pAT133;Egr4

产品验证



Western blot analysis of Egr4 expressed in Mouse brain, Mouse kidney using Egr4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn