

CXCR1 Rabbit pAb

货号: **AYP17609**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	39kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Raji
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human CXCR1 (NP_000625.1).
序列	IFLLCWLPYNLVLADTLMRTQVIQESCERRNNIGRALDATEILGFLHSLNPIIYAFIQNFRHGFLKILAMHGLVSKEFLARH RVTSYTSSSVNVSSNL

靶点信息

研究背景	The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. Knockout studies in mice suggested that this protein inhibits embryonic oligodendrocyte precursor migration in developing spinal cord. This gene, IL8RB, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36.
------	---

基因ID	3577
基因名	CXCR1
Swiss	P25024
别名	CXCR1;C-C;C-C-CKR-1;CD128;CD181;CDw128a;CKR-1;CMKAR1;IL8R1;IL8RA;IL8RBA

产品验证



Western blot analysis of CXCR1 expressed in Jurkat,Raji using CXCR1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn