

ZNF189 Rabbit pAb

货号: **AYP17641**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	62kDa/68kDa/71kDa/72kDa
实测分子量	73kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human ZNF189 (NP_001265160.1).
序列	MASPSPPESKEEWDYLDPAQRSLYKDVMMENYGNLVS LDV LNRDKDEEPTVKQEIEIEIEEVEPQGVIVTRIKSEIDQDP MGRETFELVGR LDKQ RGI FLWEIPRESLTQEQRMFRENTNII RKRPNSEEKCHKCEECGKGFVRKAHFI

靶点信息

研究背景	Kruppel-like zinc finger proteins such as ZNF189 contain a conserved stretch of 7 amino acids that connects a variable number of DNA-binding zinc finger repeats of the cys(2)his(2) (C2H2) type. Approximately 30% of human Kruppel-like zinc finger proteins contain an N-terminal Kruppel-associated box (KRAB) domain. The KRAB domain consists of approximately 75 amino acids that may be subdivided into an A box, which is present in every KRAB domain and is essential for transcriptional repression, and a B box, which is not always present.
------	--

基因ID	7743
基因名	ZNF189
Swiss	O75820
别名	ZNF189

产品验证



Western blot analysis of ZNF189 expressed in Jurkat using ZNF189 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn