

— ABLYBIO, Help Your Research



BRAP Rabbit pAb

货号: **AYP17652**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	47kDa/67kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HepG2,MCF7,mouse brain,rat testis,rat brain
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human BRAP (NP_006759.3).
------	--

靶点信息

研究背景	The protein encoded by this gene was identified by its ability to bind to the nuclear localization signal of BRCA1 and other proteins. It is a cytoplasmic protein which may regulate nuclear targeting by retaining p proteins with a nuclear localization signal in the cytoplasm.
基因ID	8315
基因名	BRAP
Swiss	Q7Z569 (https://www.uniprot.org/uniprotkb/Q7Z569/entry)
别名	BRAP, BRAP2, IMP, RNF52, BRAP Rabbit pAb, Impedes mitogenic signal propagation, RING finger protein 52, RING-type E3 ubiquitin transferase BRAP2, Renal carcinoma antigen NY-REN-63

产品验证



Western blot analysis of BRAP expressed in mouse brain, rat testis, U-251MG using BRAP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)