

# ZNF263 Rabbit pAb

货号: AYP17660

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	77kDa
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, HT-29, SW480, BxPC-3
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-240 of human ZNF263 (NP_005732.2).
序列	QRELGRLRQVQVTNHGRGTEVLLEEPLPLETARESPSFKLEPMETERSPGPRLQELLGPSPQRDPQAVKERALSAPWLSLFP PEGNMEDKEMTGPQLPESLEDVAMYISQEEWGHQDPSKRA

## 靶点信息

研究背景	Transcription factor that binds to the consensus sequence 5'-TCCTCCC-3' and acts as a transcriptional repressor. Binds to the promoter region of SIX3 and recruits other proteins involved in chromatin modification and transcriptional corepression, resulting in methylation of the promoter and transcriptional repression. Acts as transcriptional repressor of HS3ST1 and HS3ST3A1 via binding to gene promoter regions.
基因ID	10127

基因名	ZNF263
Swiss	O14978
别名	ZNF263;FPM315;ZKSCAN12;ZSCAN44

## 产品验证



Western blot analysis of ZNF263 expressed in U-87MG, HT-29, SW480 using ZNF263 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)