

DYRK4 Rabbit pAb

货号: **AYP17661**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	59kDa/72kDa/73kDa
实测分子量	73kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,Jurkat,HepG2,C2C12,Rat testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 330-480 of human DYRK4 (NP_003836.1).
序列	TASRRQTFFDKSGFKNITNNRGKKRYPDSKDLTMVLKTYDTSFLDFLRRLVWEPRLMTPDQALKHAWIHQSRNLKPQPRPQTLRKSNSFFPSETRKDKVQGGCHSSRKADEITKETTEKTKDSPTKHVQHSGDQQDCLQHGADTVQLP

靶点信息

研究背景	This gene encodes an enzyme that belongs to a conserved family of serine/threonine protein kinases. Members of this dual specificity kinase family are thought to function in the regulation of cell differentiation and proliferation, survival, and in development. Alternate splicing results in multiple transcript variants. A dditional alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known.
基因ID	8798

基因名	DYRK4
Swiss	Q9NR20
别名	DYRK4

产品验证



Western blot analysis of DYRK4 expressed in Rat testis,293T,A-549 using DYRK4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn