

— ABLYBIO, Help Your Research



COMMD5 Rabbit pAb

货号: **AYP17688**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	24kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SGC996,293T,LO2,HeLa,A-431,Mouse kidney,Rat liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-224 of human COMMD 5 (NP_054785.2).
------	---

靶点信息

研究背景	May modulate activity of cullin-RING E3 ubiquitin ligase (CRL complexes. Negatively regulates cell proliferation. Negatively regulates cell cycle G2/M phase transition probably by transactivating p21/CDKN1A through the p53/TP53-independent signaling pathway. Involved in kidney proximal tubule morphogenesis (By similarity. Down-regulates activation of NF-kappa-B.
基因ID	28991
基因名	COMMD5
Swiss	Q9GZQ3 (https://www.uniprot.org/uniprotkb/Q9GZQ3/entry)
别名	COMMD5,HCARG,HT002,COMMD5 Rabbit pAb,Hypertension-related calcium-regulated gene protein

产品验证



Western blot analysis of COMMD5 expressed in Mouse kidney,Rat liver,SGC996 using COMMD5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)