

ITSN2 Rabbit pAb

货号: **AYP17692**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	135kDa/141kDa/190kDa/193kDa
实测分子量	180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,LO2,Jurkat,HeLa
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 500-620 of human ITSN 2 (NP_006268.2).
序列	GKHQQISGRLQDVRLKKQTQKTELEVLDKQCDLEIMEIKLQQLQELQEYQNKLIYLVPEKQLLNERIKNMQFSNTPDSGVSL LHKKSLEKEELCQRLKEQLDALEKETASKLSEMSDFNNQL

靶点信息

研究背景	This gene encodes a cytoplasmic protein which contains SH3 domains. This protein is a member of a family of proteins involved in clathrin-mediated endocytosis. Intersectin 2 is thought to regulate the formation of clathrin-coated vesicles and also may function in the induction of T cell antigen receptor (TCR) endocytosis.
基因ID	50618

基因名	ITSN2
Swiss	Q9NZM3
别名	ITSN2;PRO2015;SH3D1B;SH3P18;SWA;SWAP

产品验证



Western blot analysis of ITSN2 expressed in U-87MG, LO2, Jurkat using ITSN2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn