

— ABLYBIO, Help Your Research



CYSLTR2 Rabbit pAb

货号: **AYP17717**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	mouse kidney
细胞定位	Cell membrane, Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human CYSLTR2 (NP_001295394.1).
------	---

靶点信息

研究背景	The cysteinyl leukotrienes LTC ₄ , LTD ₄ , and LTE ₄ are important mediators of human bronchial asthma. Pharmacologic studies have determined that cysteinyl leukotrienes activate at least 2 receptors, the protein encoded by this gene and CYSLTR1. This encoded receptor is a member of the superfamily of G protein-coupled receptors. It seems to play a major role in endocrine and cardiovascular systems.
基因ID	57105
基因名	CYSLTR2
Swiss	Q9NS75 (https://www.uniprot.org/uniprotkb/Q9NS75/entry)
别名	CYSLTR2,CYSLT2,CYSLT2R,GPCR21,HG57,HPN321,KPG_011,PSEC0146,hGPCR21,CYSLTR2 Rabbit pAb,G-protein coupled receptor GPCR21,G-protein coupled receptor HG57

产品验证



Western blot analysis of CYSLTR2 expressed in mouse kidney using CYSLTR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)