

ACBD3 Rabbit pAb

货号: AYP17726

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	60kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Mouse testis,Rat testis
细胞定位	Cytoplasmic side,Golgi apparatus membrane,Mitochondrion,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human ACBD3 (NP_073572.2).
序列	MAAVLNAERLEVSVDGLTLSPDPEERPGAEGAPLLPPPLPPSPPPGSRGPGASGEQPEPGEEAAGGAAEEARRLEQRWGFLEELYGLALRFFKEKDGKAFHPTYEEKLKLVALHKQVLMGPYNPDTCPVEGFFDVLGNDRRREWAALGNMSKEDAMV E

靶点信息

研究背景	The Golgi complex plays a key role in the sorting and modification of proteins exported from the endoplasmic reticulum. The protein encoded by this gene is involved in the maintenance of Golgi structure and function through its interaction with the integral membrane protein giantin. It may also be involved in the hormonal regulation of steroid formation.
------	--

基因ID	64746
基因名	ACBD3
Swiss	Q9H3P7
别名	ACBD3;GCP60;GOCAP1;GOLPH1;PAP7

产品验证



Western blot analysis of ACBD3 expressed in Mouse testis,Rat testis,HeLa using ACBD3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn