

FAM65A Rabbit pAb

货号: **AYP17728**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	131kDa/132kDa/133kDa/134kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1004-1223 of human FAM65A (NP_001180451.1).
序列	LSLRQPGLAEAVCVKFLLEDALGQKLPRRPQPGPGEQLTVFQFWSFVETLDSPTMEAYVTETAEVLLVRNLNSDDQAVVLKALRLAPEGRLRRDGLRALSSLLVHGNNKVMAAVSTQLRSLSLGPTFRERALLCFLDQLEDEDVQTRVAGCLALGCIKAPEGIEPLVYLCQTDTEAVREARQSLQCGEEGQSAHRRLEESLDALPRIFGPGSMSTAF

靶点信息

研究背景	Downstream effector protein for Rho-type small GTPases that plays a role in cell polarity and directional migration. Acts as an adapter protein, linking active Rho proteins to STK24 and STK26 kinases, and hence positively regulates Golgi reorientation in polarized cell migration upon Rho activation. Involved in the subcellular relocation of STK26 from the Golgi to cytoplasm punctae in a Rho- and PDCD10-dependent manner upon serum stimulation.
------	--

基因ID	79567
基因名	RIPOR1
Swiss	Q6ZS17
别名	RIPOR1;FAM65A

产品验证



Western blot analysis of FAM65A expressed in Mouse kidney using FAM65A Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn