

KIF17 Rabbit pAb

货号: **AYP17731**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	114kDa/115kDa
实测分子量	170kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,U-87MG,PC-3,Mouse brain,Mouse testis
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 360-660 of human KIF17 (NP_001116291).
序列	LKAILTQQMSPSSLSALLSRQVPPDPVQVEEKLLPQPVIQHDVVEAEKQLIREEYEERLARLKADYKAEQESRARLEEDITAMR NSYDVRSLSTEENLRKETEAVLQVGVLYKAEVMSRAEFASSAEYPPAFQYETVVKPKVFSTDTLPSDDVSKTQVSSRFAE LPKVEPSKSEISLGSSSESSLEETSVSEAFPGPEEPSNVEVSMPTTEESRSRYFLDECLGQEAAGHLLGEQNYLPQEPEVP LQGLLGLQDPFAEVEAKLARLSSTVARTDAPQADVPKVPVQVPAPTDLLEPSDAR

靶点信息

研究背景	Dendrite-specific motor protein which, in association with the Apba1-containing complex (LIN-10-LIN-2-LIN-7 complex, transports vesicles containing N-methyl-D-aspartate (NMDA receptor subunit NR2B along microtubules.
------	--

基因ID	57576
基因名	KIF17
Swiss	Q9P2E2
别名	KIF17;KIF17B;KIF3X;KLP-2;OSM-3

产品验证



Western blot analysis of KIF17 expressed in Mouse brain, HeLa, U-87MG using KIF17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn